

.Sikdar S, Sadhukhan S, Majumdar AK, Bhunia S, Sarkar S, Ganguly S. Phenotypic Characterisation, Virulence Determination and Antimicrobial Resistance Pattern of Enterococcus Species Isolated from Clinical Specimen in a Tertiary Care Hospital in Kolkata. Journal of Clinical and Diagnostic Research, 2021; 15(7): DC06-DC09.Sikdar S, Bhunia S, Majumdar AK, Sarkar S, Ganguly S Inconclusive in COVID-19 Reverse Transcriptase-Polymerase Chain Reaction Test: A Diagnostic Dilemma. Int J Cur Res Rev, 2021; 13 (11): S105 – 108.Deb S, Choudhury K, Sarkar S, Mukherjee S, Ganguly S, Pal NK. Bacteriocin production by clinical isolates of ‘Non fermenting Gram negative bacilli’ – a ray of hope in the post antibiotic era. International Journal of Health and Clinical Research, 2021;4(6):55 – 58. Dutta S, Choudhury K, Banerjee C, Ghosh A, Sarkar S, Ganguly S. Study on parameters of severity in Dengue infection with both NS1 antigen and Dengue IgM reactive on the same day in a tertiary care hospital, Kolkata. International Journal of Health and Clinical Research, 2021; 4(5):84 - 87. Choudhury K, SkMd Imran Ali, Deb S, Ganguly S. Susceptibility pattern of Methicillin resistant Staphylococcus aureus (MRSA) among all clinical isolates in a tertiary care hospital in Eastern India. International Journal of Research and Review, 2020; 7(10): 546 – 550. . Deb S, Sarkar S, Chatterjee C, Ganguly S, Pal NK. Study of Gram negative non-fermenting bacilli from surgical site infections in a tertiary care hospital in Kolkata. International Journal of Research and Review, 2019; 6(7):169-174. Saha P, Ballav S, Chatterjee M, Ganguly S, Sarker M, MajiAKet al. Susceptibility status of Japanese Encephalitis vectors to insecticides in endemic areas of northern districts of West Bengal, India. Japanese Journal of Infectious Disease, 2018;71, 91–98. Saha P, Ganguly S, Chatterjee M, Das SB, Kundu PK, MajiAKet al. Asymptomatic leishmaniasis in kala-azar endemic areas of Malda district, West Bengal, India. PLoSNegl Trop Dis, 2017; 11 (2). Chatterjee M , Ganguly S, Saha P, Guha SK, Maji AK. Polymorphisms in pfdhfr and pfdhps genes after five years of artemisinin combination therapy (ACT) implementation from urban Kolkata, India. Infection, Genetics and Evolution, 2017; 52: 155–159. Guha SK, Das SB, Chatterjee M, Saha P, Ganguly S, Chakraborty A, Ganguly N and Maji AK. Epidemiological studies on leishmaniasis in Murshidabad district of West Bengal, India. International Journal of Recent Scientific Research, 2017; 8: 21660-21664. Chatterjee M, Ganguly S, Saha P, Guha SK, Basu N, Bera DK, Maji AK. Polymorphisms in Pfcr and Pfmdr-1 genes after five years withdrawal of chloroquine for treatment of Plasmodium falciparum malaria in West Bengal, India. Infection, Genetics and Evolution, 2016; 44:281–285. Saha P, Ganguly S, Maji AK Genetic diversity and multiplicity of infection of Plasmodium falciparum isolates from Kolkata, West Bengal, India, Infection, Genetics and Evolution, 2016; 43:239-44. Chatterjee M, Ganguly S, Saha P, Bera DK, Basu N, and K. Maji AK. Prevalence of G6PD Deficiency among the Malaria Patients in Urban Kolkata, India - A Clinic Based Study. Int.J.Curr.Microbiol.App.Sci. 2016; 5(3): 738-744. Ganguly S, Saha P, Chatterjee M, Roy S, Ghosh TK, MajiAKet al. PKDL—A Silent Parasite Pool for Transmission of Leishmaniasis in Kalaazar Endemic Areas of Malda District, West Bengal, India. PLoSNegl Trop Dis, 2015; 9(10): e0004138. doi:10.1371/journal.pntd.0004138. Chatterjee M, Ganguly S, Saha P, Bankura B, Basu N, Das M, Guha SK, and Maji AK. No Polymorphism in Plasmodium falciparum K13 Propeller Gene in Clinical Isolates from Kolkata, India. Journal of Pathogens, Volume 2015, Article ID 374354, <http://dx.doi.org/10.1155/2015/374354>. Saha P, Ganguly S, Saha P, Chatterjee M, Kundu PK and MajiAK.Comparison of three diagnostic methods for the detection and speciation of malarial parasite in urban Kolkata, India, International Journal of Parasitology Research, 2015; 7: 160-163Ganguly S, Saha P, Chatterjee M and Maji AK. Prevalence of polymorphisms in antifolate drug resistant molecular marker genes, pvdhfr and pvdhps in clinical isolates from Kolkata, India, Antimicrobial Agent and Chemotherapy, 2014; 58(1):196-200. DOI: 10.1128/AAC.01171-13. Ganguly S, SahaP,Guha SK, Basu N, and Maji AK. Recurrence pattern of P vivax malaria following treatment

with chloroquine either alone or in combination with Primaquine in urban Kolkata, India,
International Journal of Recent Scientific Research, 2014;5: 1046 -1049